

Variance of some Soil Physicochemical Properties and Oxides within the Oases of Al-Bouhyat in the Anbar Desert, Western Iraq

Mohammed A. Al-Fahdawi¹, Salah M. Al-Juraysi^{2*}

¹ Res., Anbar Agriculture Directorate, Ministry of Agriculture, Anbar, Iraq.

² Prof., Department of Soil and Water Resources Science, College of Agriculture, University and Anbar, Anbar, Iraq.

Abstract

Al-Bouhyat Oasis was chosen. It is located within the boundaries of Anbar Governorate, to the right of the Euphrates River, southeast of Lake Haditha, about 27.45 km, at the coordinates 33°57'59.4"N 42°26'25.4"E, and an area of about 146.55 ha., Soil survey was carried out using the Free Lance method, covered with several pedons with a semi-detailed survey, which numbered 6 pedons. Soil samples were taken from each diagnosed horizon, transferred to the laboratory, and prepared for some physical and chemical analyses. The results showed soil development at the calcareous materials. The soils were classified into the Order of Aridisols, which included two levels of the suborder Gypsid with 78.25 ha (53.39%) and Calcids with 68.30 hectare (46.61%). The study results showed the SiO₂ content in all profiles decreased with depth (range of 28.98 % to 35.14 and that Al₂O₃ values varied from 5.3 to 8.65%. The highest amount of Fe₂O₃ (4.02%) was in series 131CCW. The CaO range from 21.87 to 31.8% in series 133CKW and series 122CCW. The correlation coefficients showed no significant correlations with depth for Fe₂O₃, SiO₂ and Al₂O₃, with a correlation coefficient of -0.096, 0.093 and 0.086, respectively. The study also showed a positive correlation coefficient between gypsum and sand content, with a significant positive correlation coefficient of 0.67. This indicates the coarse texture of gypsum soils. The study also showed a significant negative correlation coefficient of -0.51 between the gypsum content and the bulk density.

Keywords: Gypsid, elemental oxide, calcium oxide, SPSS

تباين بعض الصفات الفيزيوكيميائية للتربة وأكاسيدها ضمن واحة البوحيات في صحراء الأنبار غربي العراق

محمد أحمد الفهداوي¹, صلاح مرشد الجريسي^{2*}

¹ باحث، مديرية زراعة الأنبار، وزارة الزراعة، الأنبار، العراق.

² استاذ، قسم علوم التربة والموارد المائية، كلية الزراعة، جامعة الأنبار، الأنبار، العراق.

المستخلص

اختيرت واحة البوحيات والتي تقع ضمن حدود محافظة الأنبار، على يمين نهر الفرات جنوب شرق بحيرة حديثة بحوالي 27.45 كم، تقع عند الإحداثيات الجغرافية 33°57'59.4" شرقاً و 42°26'25.4" شمالاً. وتبلغ مساحتها 146.55 هكتار. تم إجراء مسح للتربة باستخدام الطريقة الحرة Free Lance حفرت 6 بيدونات ممثلة للتربة المتوقع وجودها. استحصلت عينات التربة من افاق البيدونات وحددت حدود الافاق وسمك الافاق ونقلت النماذج الترابية إلى المختبر لإجراء بعض التحليلات الفيزيائية والكيميائية والاكاسيد المهمة في تصنيف التربة حسب النظام الأمريكي Soil Survey Staff. 2014 ونظام تصنيف العكدي للتربة الصحراوية 1981. أظهرت النتائج أن التربة نشأت من مواد اصل كلسية. وصنفت التربة عند مستوى رتبة Aridisols والتي صنفت الى مستويين من تحت الرتبة Gypsid وبمساحة 78.25 هكتار ونسبة 53.39% والمستوى تحت الرتبة Calcids وبمساحة 68.30 هكتار ونسبة 46.61%. لوحظ من خلال النتائج المتحصل عليها انخفاض في نسبة اوكسيد السليكون SiO₂ في جميع البيدونات مع العمق من 35.14% الى 28.98%، كما لوحظ أن نسبة اوكسيد الالمنيوم Al₂O₃ تباينت من 5.3 إلى 8.65%. وكانت أعلى قيمة لاوكسيد الحديد Fe₂O₃ 4.02% في السلسلة 131 CCW اما اوكسيد الكالسيوم CaO فقد تراوح ما بين 21.78 إلى 31.8% في السلسلة 133CKW والسلسلة 122CCW بالتتابع. اوضحت معاملات الارتباط عدم وجود علاقة ارتباط مع عمق التربة لاكاسيد الحديد والسليكون والالمنيوم وبمعامل ارتباط -0.096، 0.093 و 0.086 على التوالي. كما بينت الدراسة وجود معامل ارتباط موجب بين محتوى الجبس ومحتوى الرمل وبمعامل ارتباط 0.67. وهذا يشير الى خشونة نسجة التربة الجبسية. كما اوضحت الدراسة وجود معامل ارتباط سالب بين محتوى الجبس والكثافة الظاهرية وبمعامل ارتباط -0.51.

الكلمات المفتاحية: واحة البوحيات، اوكسيدات العناصر، اوكسيد الكالسيوم، SPSS

*Corresponding author.

Email: ag.salah.murshid@uoanbar.edu.iq

https://dx.doi.org/10.36531/ijds.2022.176697

Received 30 August 2022; Received in revised form 27 September 2022; Accepted 9 October 2022

Introduction

The particle size distribution as a fundamental factor generally controls spatial variations of soil physical properties in the field. Heterogeneity and variation of soil physical parameters in a field due to soil plowing should be considered for successful agricultural management (Gulser et al., 2016).

Soil physical and chemical properties are strongly influenced by soil management systems and changes in land use (Hulugalle et al., 1997).

The main aim of statistical testing is to attach probabilities to those inferences. The tests are prefaced by hypotheses or models, such as no differences among population means or relations between variables, the so-called null hypotheses (Webster, 2021). A typical geostatistical analysis of soil data proceeds on the assumption that the properties of interest are the outcomes of random processes. Many factors have contributed to the soil in the parent material and during its formation (Webster, 2000).

Mzuku et al. (2005) showed that soil physical properties exhibited significant spatial variability across production fields. The trends observed for the measured soil physical properties correspond to the productivity potential of the management zones. Utilizing site-specific management zones could help manage the in-field variability of yield-limiting soil physical properties.

Al-Juraysi, (2021) Mention There is no apparent chemical weathering in the study pedons and for all indications, due to the little leaching of calcium carbonate in sufficient quantities due to the little rain and the

accumulation of quartz in the soil surface, which indicates that the soils are newly formed.

Al-Khafagi et al. (2016) showed an apparent increase in immobile oxides of (SiO_2 , Fe_2O_3 and Al_2O_3). However, the mobile oxides of (CaO) and the loss in the ignition were increased in Al-Hadher soil and gradually decreased in Al-Qayarah, Hamam Al-All, and Tallafer and reached their lowest levels in Faidah soil. Research on soil variation could be improved by increased coordination of experimental designs and by application of statistical methods suitable for analysis of place-to-place variation, including statistical functions such as spatial autocorrelation (Campbel, 1979).

In a study conducted by Beckett and Webster, 1971 They showed that the variations in soil characteristics from one place to another in the terrestrial perspective are the result of several reasons, including the climate, which causes gradual and sudden changes; topography and material of origin, which cause local differences and short-term variances; and vital activities that cause regional differences.

The state of weathering, its intensity and direction can be known by studying the mineral composition of the soil and the percentage of oxides in the soil volume. He added that the evidence of soil weathering, which is based on the proportions and quantity of oxides and the proportions of minerals present, is essential in the study of the evolutionary state of the soil. (Aljoboury, 2018).

The differences in soil formation factors and processes are due to variables such as parent material, topographical position, and age of

the associated landscape (Dengiz and Usul, 2018).

This research aims to detect variations in some physical and chemical properties and oxides in dry desert soils.

Materials and Methods

The study area is located at coordinate central 33 °57'59.4"N 42°26'25.4"E, within the boundaries of Anbar Governorate, shown in Figure 1, which is located to the right of the Euphrates River, southeast of Lake Haditha about 27.45 km, this study was carried out on an area of approximately 146.55 ha. Soil samples were collected from six representative profiles in the Al-Bouhyat Oasis through a semi-detailed survey. The coordinates of the pedons' locations were recorded using the GPS device GARMIN. After the soil samples taken from six soil profiles were air-dried, each sample was passed out of a 2 mm sieve. Particle size distribution was determined with the hydrometer method (Bouyoucos, 1962). Bulk density was determined by using undisturbed samples (Black, 1965). and cation exchange capacity (CEC) was measured with the 1 N NH₄OAc (pH7) method (Soil Survey Laboratory Manuals, 2004).

The oxides were analyzed in the laboratory of chemical and mineral analyzes of the Fallujah Cement Factory using an X-ray Fluorescence (XRF) device. The climate is an arid continental climate, and its annual precipitation is 144.85 mm. The mean annual temperature is about 20.36°C. Different types, including desert, Haloxylon

salicornicum, Guisum Achillee and Atriplex spp, characterize the natural vegetation of the study area. The soils were classified to the family level based on (Soil Survey Staff, 2014).

Statistical analysis

The relationships between the soil properties were evaluated by obtaining the values of Pearson's correlation, and Descriptive statistics of the soil properties (Mean, Std., Deviation, Minimum, Maximum and coefficient of variation) were calculated using the SPSS program for selected soil properties.

Results and Discussions

The major physical, chemical, and oxide characteristics of the soils in each pedon in the study area are presented in Table 1. Physical and chemical characteristics in the different pedons varied due to a dynamic interaction between climate and parent material (Kibar *et al.*, 2012). Solum depth ranges from 30 to 100 cm, depending upon the stage of the soil formation process. the delineation of soil horizons is affected by lateral and vertical variation in soil properties, which are texture, gypsum, lime, ECe, ESP, and CEC as their importance increases at the soil surface and decreases with depth. Al-Juraysi and Mhede (2021). The dominant soil texture for pedons 2 and 4 was loam and sandy loam. The highest sand content was 864 g kg⁻¹ in series 131CCW pedon1, formed on Calcareous and gypsiferous material parent material.




Figure 1. The location of the study area

Gypsum soils generally contain no more than 15% clay. Soil texture largely depends on the nature of the parent material from which the soil is formed and the degree of overlap in gypsum deposits. Very high calcium carbonate content was noted in series 131CCW pedon3 and series 122CCW Pedon5 (485 and 495 g kg⁻¹) respectively (Table.1). The original parent materials provided the necessary calcium. This explanation is plausible for parent materials containing limestone and other calcium-rich rocks. While In series 131CCW Pedon1(A) the content of calcium carbonate was 100 g kg⁻¹, while series 132XKM pedon4(By) only 290 g kg⁻¹. (Table 1). The content and distribution of the gypsum was noted in series 131CCW pedon 1, Pedon3, pedon4 and pedon6 ,340, 223, 280 and 223 g kg⁻¹, respectively. Soil bulk density values ranged from 1.15 to 1.75 g cm⁻³. with values generally higher in the

subsurface horizons as a result of compaction. The low bulk density is due to a relatively low specific weight of gypsum, e.g., 2.3 gr per cm³, in combination with an occasional high porosity. Soil Cation Exchange Capacity varied from 10.5 to 32.8 cmolc kg⁻¹, with the highest CEC in the series 133CKW p6 horizons. The lowest CEC value was found in soil classified as series 131CCW pedon p1 horizons C_{K2}. This capacity is about inversely proportional to the gypsum content. When the gypsum content increases, the amount of the non-gypsiferous components, including clay, decreases. The ratio concentrations of mobile and immobile elements.

Table 2. shows that the SiO₂ content in all profiles decreased with a depth range of 28.98 to 35.14 % in series 133CKW pedon 6 horizon C_{K1} and series 122CCW pedon 5 horizon A_K respectively. that Al₂O₃ values

varied from 5.3 to 8.65% in series133CKW pedon p6 horizon C_{K1} and series132XKM pedon p4 horizon A_K respectively. SiO₂ highly resists weathering because it is contained in quartz minerals. In contrast, Al₂O₃ is less resistant to weathering because it is contained in clay minerals; a high content of Al₂O₃ indicates a high content of clay minerals (Demiralay, 1993; Shan et al., 2010). clay152 and 154 g.kg⁻¹, series132XKM pedon p4 horizon A_K and

series 122CCW pedon p5 horizon A_K. respectively; Table.2. The highest amount of Fe₂O₃ 4.02% was in series131CCW pedon p1. Concentrations of CaO are seen in the soils where the parent materials are mostly limestone. Generally, the CaO content in these soils ranges from 21.87 to 31.8% in series 133CKW pedon p6 horizon C_{K1} and series 122CCW pedon5 horizon B_K, respectively.

Table 1. Some physical and chemical characteristics of soil in the study area

pedon	Horizon	Depth / cm	ECe	Lime gkg ⁻¹	Gypsu m gkg ⁻¹	CEC cmolc kg ⁻¹	Sand gkg ⁻¹	clay gkg ⁻¹	silt gkg ⁻¹	texture	bulk density gcm ³
P1	A	0-16	0.5	100	140	19.9	428.0	96.0	476.0	L	1.46
	B _{yk}	16-29	4.2	275	340	16.8	763.0	97.0	140.0	SL	1.32
	C _{K1}	29-60	13.0	375	280	18.3	864.0	67.0	69.0	LS	1.15
	C _{K2}	60-100	14.0	400	220	10.5	764.0	68.0	168.0	SL	1.38
p2	A	0-15	1.8	350	150	10.8	387.0	137.0	476.0	L	1.57
	B _K	15-30	3.5	465	160	14.9	763.0	112.0	125.0	SL	1.60
P3	A	0-11	32.2	485	Nil	17.0	337.0	161.0	502.0	SiL	1.39
	B _y	11-24	19.4	485	223	17.3	634.0	121.0	245.0	SL	1.39
	C _{K1}	24-64	19.8	485	50	11.3	703.0	92.0	205.0	SL	1.33
	C _{K2}	64-90	15.1	485	165	13.9	664.0	98.0	238.0	SL	1.40
P4	A _K	0-22	9.0	410	140	18.9	376.0	151.0	473.0	L	1.21
	B _y	22-48	6.7	290	280	12.0	772.0	105.0	123.0	SL	1.25
P5	A _K	0-48	2.7	475	Nil	17.4	545.0	154.0	301.0	SL	1.56
	B _K	48-61	3.4	490	Nil	17.2	196.0	306.0	498.0	SiCL	1.61
	C _{K1}	61-81	2.6	495	Nil	12.0	225.0	253.0	522.0	SiL	1.75
P6	A _K	0-27	0.9	470	Nil	14.8	373.0	176.0	451.0	L	1.39
	B _K	27-57	2.1	450	15	22.0	667.0	130.0	203.0	SL	1.58
	B _y	57-97	6.6	325	223	32.8	518.0	278.0	204.0	SCL	1.61

Table 2. The elemental oxide concentrations of soils in the study area

Pedon No	Horizon	Depth (cm)	Na ₂ O	K ₂ O	SiO ₂	MgO	CaO	Fe ₂ O ₃	Al ₂ O ₃
P1	A	0-16	0.11	1.18	32.23	4.612	24.06	4.02	6.74
	B _{yk}	16-29	0.13	0.48	30.04	4.558	24.07	3.27	5.89
	C _{K1}	29-60	0.13	0.57	30.96	4.744	24.45	3.10	6.88
	C _{K2}	60-100	0.15	0.76	30.45	4.677	24.92	3.10	6.19
P2	A _K	0-15	0.12	0.16	31.69	6.614	25.03	3.78	6.98
	B _K	15-30	0.1	0.77	31.31	4.749	27.7	3.17	6.68
P3	A _K	0-11	0.32	1.25	31.63	4.661	25.3	3.65	6.87
	B _K	11-24	0.2	0.78	30.84	4.724	26.31	3.05	6.57
	C _{K1}	24-64	0.19	0.8	31.45	4.883	29.07	2.68	6.95
	C _{K2}	64-90	0.26	0.26	30.46	4.723	27.43	2.67	6.65
P4	A _K	0-22	0.17	1.09	34.4	4.784	26.06	3.77	8.65
	B _K	22-48	0.12	0.47	30.79	4.777	26.57	2.79	6.60
P5	A _K	0-48	0.09	1.27	35.14	4.763	25.83	3.77	8.63
	B _K	48-61	0.1	0.82	32.28	4.916	31.8	2.87	7.57
	C _K	61-81	0.12	1.03	30.68	4.776	31.03	2.81	6.39
P6	A _K	0-27	0.12	1.89	31.96	4.579	24.43	3.77	6.86
	B _K	27-57	0.11	1.03	31.08	4.79	25.49	3.05	6.97
	B _y	57-97	0.12	1.52	28.98	4.501	21.87	3.39	5.30

Soil classification of the study area

The region's soils were classified according to the modern american classification (Table 3.) into the order of aridisols, which included two levels of the suborders Gypsid with 78.25 hectare 53.39% with two levels of the family, and Calcids with 68.30 hectare 46.61%. With one level of the family coarse loamy, mixed, hyperthermic, super active, Typic petrogypsid. It is a level of desert soil with a moderately coarse texture, originating from materials of limestone origin that have a petroglyphic horizon within 100 cm of the soil surface. (Soil Survey Staff., 2014) Parent material is calcareous and gypsiferous material. The distribution of lime decreases in the second horizon. While gypsum increased with the depth, Flat lands with low slopes of 1% occupied an area of 22.31 hectares, or 15.22% of the total area of the

oasis, the natural plant available were *achiiea fragrantissima*, *piatynerium bifurcatum* and *urtica dioica*. It was represented by the pedon 4.

Coarse- loamy, mixed, hyperthermic, super active, Typic calcigypsid. Gypsid that have a calcic horizon within 100 cm of the soil surface. (Soil Survey Staff., 2014) with a moderately coarse texture originating from calcareous materials. The texture of its horizons is characterized by its mixture in the upper horizon. Sandy mixture in the rest of the horizons, in which the distribution of lime appears regularly, as it increases with increasing depth. The lands of this series are located within flat lands with a slope of 1% and occupied an area of 55.94 hectares or 38.17% of the total area of the oasis (Table 4) in which *Haloxylon Salicornicum*, *artemisia herba alba* and *haioxion*. represented by the

pedon 1, pedon 3 and pedon 6. (Figure 2) Coarse loamy, mixed, hyperthermic, super active, Typic Haplocalcids.

It is a level desert soil with a moderately coarse texture and originated from limestone materials. the distribution of lime appears

regularly with the depth. Within the flat lands with a slight slope ranging between (1-2%), it occupied an area of 68.30 hectares, with a percentage of 46.61% of the total area of the oasis. (Table 4) represented by pedon 2 and pedon 5 (Figure 2)

Table 3. Soil classification of the study area

order	Sub order	Great group	Sub great group	Family	series	pedons
Aridisols	Gypsisols	Petrogypsisols	Typic Petrogypsisols	Coarse loamy, mixed, hyperthermic, super active topic petrogypsisols	132XKM	4
					131CCW	1,3
					133CKW	6
	Calcids	Haplocalcids	Typic Haplocalcids	Coarse loamy, mixed, hyperthermic, super active, Typic calcigypsisols	132CCM	2
					122CCW	5

Table 4. Area of Soil classification of the study area

No.	Sub order	series name	Area/hectare	Percentage (%)	pedons
1	Gypsisols	G3 131 CCW	33.49	22.85	1,3
		G3 133 CKW	22.45	15.32	6
		G2 132 XKM	22.31	15.22	4
		Sum	78.25	53.39	
2	Calcids	G2 132 CCM	50.88	34.72	2
		G3 122 CCW	17.42	11.89	5
		Sum.	68.30	46.61	
the total area			146.55	100	




Figure 2. The soil series and locations of the selected pedons Statistical analysis: correlation coefficient

the correlations among the soil properties were calculated using the SPSS program. Most soil properties were significantly positively and negatively correlated with each other. The table.5 shows the correlation among soil properties. All soil properties had no significant correlations with depth. Silicon dioxide (SiO₂) content in all profiles decreased with depth. It strongly resists weathering because it is mainly contained in quartz minerals. With a positive, no significant correlation coefficient of 0.093. Aluminium oxide (Al₂O₃) content in all profiles decreased with depth. It is mainly less resistant to weathering because it contains clay minerals. High content of Al₂O₃ indicates an increased range of clay minerals. (Shan et al., 2010). with a positive correlation coefficient of .086. Iron oxide (Fe₂O₃) content in all profiles decreased with depth. With a negative correlation coefficient of -.096. Potassium Oxide (K₂O) had significant positive correlations with Fe₂O₃ 0.47 and SiO₂ 0.25 and significant negative correlations with MgO -0.49. SiO₂ had significant positive correlations with Al₂O₃ 0.96 and Fe₂O₃ 0.51. Calcium Oxide (CaO) had significant negative correlations with Fe₂O₃ -

0.58. This situation can be explained by the modifications occurring due to the effects of environmental factors over time . The electrical conductivity (ECe) had significant positive correlations with Na₂O 0.89 and lime content 0.34. and significant negative correlations with bulk density -0.43, clay -0.25, Fe₂O₃ -0.22 and MgO -0.18. Lime had significant negative correlations with Gypsum -0.54 and Fe₂O₃ -0.39 and significant positive correlations with CaO 0.55. Gypsum had significant negative correlations with silt content -0.65, K₂O -0.52, bulk density -0.51, SiO₂ -0.49, Al₂O₃ -0.47, CaO -0.46 and clay -0.34. The poor K status of soils containing a high percentage of Gypsum and significant positive correlations with Sand content is 0.67 because a gypsic layer can have either a powdery or a sandy appearance, depending on the size of the gypsum crystals, which may vary from 50 to over 2,000 microns. Soil Cations Exchange Capacity content (CEC) had significant negative correlations with CaO -0.49 and MgO -0.37 and significant positive correlations with clay content 0.40 and K₂O 0.49.

Table 5. Pearson’s correlation coefficient values show the relationship among the study area soil properties.

properties	Depth cm	SiO ₂ %	MgO%	CaO%	Na ₂ O%	K ₂ O%	Fe ₂ O ₃ %	Al ₂ O ₃ %	ECe dSm ⁻¹	Lime gkg ⁻¹	Gypsum gkg ⁻¹	CEC cmolc kg ⁻¹	Sand g.kg ⁻¹	silt g.kg ⁻¹	clay g.kg ⁻¹	density
Depth cm	1															
SiO ₂ %	.093	1														
MgO%	-.205	.126	1													
CaO%	-.215	.171	.078	1												
Na ₂ O%	-.207	-.126	-.122	.008	1											
K ₂ O%	.250	.25	-.049*	-.235	-.105	1										
Fe ₂ O ₃ %	-.096	.51*	.170	-.589*	-.115	.47*	1									
Al ₂ O ₃ %	.086	.96**	.168	.293	-.053	.097	.310	1								
ECe dSm ⁻¹	-.034	-.157	-.180	.008	.894**	-.104	-.222	-.076	1							
Lime gkg ⁻¹	.116	.151	-.015	.548*	.320	.094	-.396	.293	.345	1						
Gypsum gkg ⁻¹	-.064	-.48*	-.052	-.462	-.058	-.518*	-.138	-.473*	.037	-.544*	1					
CECcmolc kg ⁻¹	.151	-.136	-.368	-.493*	-.134	.485*	.277	-.172	-.116	-.247	.083	1				
Sand gkg ⁻¹	.345	-.356	-.214	-.321	-.025	-.431	-.371	-.292	.164	-.157	.669**	-.075	1			
Silt gkg ⁻¹	-.401	.450	.281	.271	.119	.360	.472*	.372	-.101	.087	-.65**	-.077	-.95**	1		
clay gkg ⁻¹	-.131	.041	-.007	.324	-.125	.380	.056	-.005	-.25	.205	-.341	.273	-.73**	.54*	1	
bulk density gcm ⁻³	-.059	-.062	.205	.319	-.345	.209	-.008	-.127	-.432	.214	-.51*	.157	-.49*	.357	.502*	1

**Correlation is significant at the 0.01 level (2-tailed) . * Correlation is significant at the 0.05 level (2-tailed)

Sand had significant negative correlations with silt content -0.96, clay -0.75, bulk density -0.49, K_2O -0.43, Fe_2O_3 -0.37 and significant positive correlations with Gypsum 0.67. Clay had significant positive correlations with a bulk density of 0.63, silt content of 0.54, and K_2O is 0.45. Silt had significant positive correlations with Fe_2O_3 0.47 and SiO_2 0.45.

Bulk density reflects the soil's ability to function for structural support, water and solute movement, and soil aeration. It had significant negative correlations with Sand -0.49 and Gypsum -0.51. The low bulk density is due to gypsum's relatively low specific weight. Significant positive correlations with clay were 0.63 and silt 0.36. It is known that the variation in bulk densities results from differences in soil texture. The parent materials are mainly limestone, and concentrations of CaO are seen in the soils. had significant positive correlations with lime 0.55. The CaO content in these soils ranged from 100 to 495 $g\ kg^{-1}$.

Statistical Analysis: Descriptive statistics

Descriptive statistics of the soil properties were calculated using the SPSS program for selected soil properties from all study sites are presented in Table 6. Mean, Std. Deviation, Minimum, Maximum and coefficient of variation, Soil properties having a coefficient of variation (CV) between 0 and 15 % are considered least variable, 15 and 35 %, moderately variable, and bigger than 35 % highly variable (Ogunkunle, 1993). Overall, the Spatial variability of soil physical properties of the sites used in this study was high in sand content, with mean sand ranging from 196.0 to 864.0 $g\ kg^{-1}$, and highly variable with a coefficient of variation of

37.2%. The silt was the most variable soil separate, ranging from 69.0 to 522.0 $g\ kg^{-1}$ and highly variable with a coefficient of variation of 53.2%. The sites used in this study were low in clay content, with a mean ranging from 13.0 to 306.0 $g\ kg^{-1}$, and highly variable with a coefficient of variation of 55.1%. The sites used in this study were low in bulk density content, with a mean ranging from 1.15 to 1.75 $g\ cm^3$, and the least variable with a coefficient of variation of 11.2%. Spatial variability of soil chemical properties of the sites used in this study was high in The electrical conductivity (ECe), ranging from 0.5 to 32.2 dSm^{-1} , and highly variable with a coefficient of variation of 98.3%. Lime content with mean ranging from 10.0 to 49.5 $g\ kg^{-1}$, and moderately variable with a coefficient of variation of 26.0%. Gypsum content with mean ranging from 0.0 to 34.0 $g\ kg^{-1}$, and highly variable with a coefficient of variation 86.1%. Soil Cations Exchange Capacity content (CEC) with mean ranging from 10.5 to 32.8 $cmolc\ kg^{-1}$, and moderately variable with a coefficient of variation of 31.7%. Distributions of all the studied variables of oxide were the least variable, and their means having a coefficient of variation (CV) between 0 and 15 %, except Na_2O and K_2O , which had CV 41.0 and 48.9%, respectively.

Distributions of all the studied variables were lightly skewed (skewness <1) and their means were close to their medians, except Na_2O , SiO_2 , MgO , ECe and CEC, which skewed with a value of 1.8, 1.2, 3.9, 1.4 and 1.7, respectively.

Table 6. properties mean and coefficient of variation (CV)for selected soil properties

properties	Descriptive statistics							
	Min.	Max.	Mean	Std. Deviation	c.v%	C. V. class	Skewness	Kurtosis
Sand g.kg ⁻¹	196.0	864.0	554.4	206.0	37.2	+++	-0.3	-1.2
silt g.kg ⁻¹	69.0	522.0	301.1	160.3	53.2	+++	0.2	-1.7
clay g.kg ⁻¹	13.0	306.0	138.1	76.0	55.1	+++	0.9	0.6
bulk density gcm ⁻³	1.15	1.75	1.4	0.2	11.2	+	0.0	-0.7
ECe dSm ⁻¹	0.5	32.2	8.7	8.6	98.3	+++	1.4	1.8
Lime gkg ⁻¹	10.0	49.5	40.6	10.5	26.0	++	-1.6	2.9
Gypsum gkg ⁻¹	0.0	34.0	13.3	11.4	86.1	+++	0.2	-1.3
CEC cmolc kg ⁻¹	10.5	32.8	16.5	5.2	31.7	++	1.7	4.8
Na ₂ O	0.1	0.3	0.1	0.1	41.0	+++	1.8	3.1
K ₂ O	0.2	1.9	0.9	0.4	48.9	+++	0.4	0.3
SiO ₂	29.0	35.1	31.5	1.5	4.6	+	1.2	2.1
MgO	4.5	6.6	4.8	0.5	9.5	+	3.9	15.7
CaO	21.9	31.8	26.2	2.5	9.5	+	0.9	0.8
Fe ₂ O ₃	2.7	4.0	3.3	0.4	13.3	+	0.3	-1.3
Al ₂ O ₃	5.3	8.7	6.9	0.8	11.8	+	0.8	1.8

C.V%: coefficient of variation, + least variable, ++ moderately variable, +++ highly variable

Conclusion

The soil of the study area belongs to the zone of arid soils, and It is located at the same soil-forming factors, topography and climate. The soil of the study area was classified into five key soils, By using the Soil Taxonomy classification system. which are differentiated in terms of their areas and their geomorphological location. The parent materials are mostly limestone.

References

Aljoboury, M. A. J. (2018). determination of homogeneity of parent material and the development of soils at selected areas in northern Iraq using weathering resistant minerals. *Iraqi National Journal of Earth Science*, 18(2), 15-34.

Acknowledgements

We would like to thank collage of agriculture/ university of Anbar for their support in conducting some analyses on the laboratory side. and laboratory of chemical and mineral analyzes of the Fallujah Cement Factory for using an X-ray Fluorescence (XRF) device contribution during the oxides analyzed.

Al-Juraysi, S. M. F. (2021). Variability of chemical weathering indices applied to desert soils formed on calcareous parent materials of western Anbar. *International Journal of Agricultural and Statistical Sciences*. 17(1) 321-327.

- Al-Juraysi, S. M & Mhede I. R. (2021). Proposing A table for weighing the characteristics of stratified samples instead of mathematical equations to study soil characteristics Statistically. *IOP Conference Series: Earth and Environmental Science*, 761.
- Al-Khafagi, Q. D., Al-Dulaimi, A. A., Esmael, A. S., & Farhan, M. J. (2016). Soil content of some oxides as a function of the development in ninavah governorate/ North Iraq. *Tikrit Journal of Pure Science*. 21(5), 139-147.
- Black, G. R. (1965). Bulk density. In Black, C. A. *et al.* (eds.) *Methods of soil analysis*, Agron. Mono. No. 9: 374-390. American Society of Agronomy, Madison, W I. USA
- Bouyoucos, G. J. (1962). Hydrometer method improved for making particle size analyses of soils. *Agronomy Journal*, 54 (5), 419-434.
- Campbell, J. B. (1979). Spatial variability of Soils. *Annals of the Association of American Geographers*. 69(4), 544–556.
- Demiralay, I. (1993). *Methods of soil physical analyses*. Agricultural Faculty, Ataturk University, Erzurum, Turkey.
- Dengiz, O., & Usul, M. (2018). Multi-criteria approach with linear combination technique and analytical hierarchy process in land evaluation studies. *Eurasian Journal of Soil Science*, 7(1), 20-29.
- Gülser, C., Imanverdi, E., Feride, C. & Zeynep, D. (2016). Spatial variability of soil physical properties in a cultivated field. *Eurasian Journal of Soil Science*, 5 (3), 192 – 200.
- Hulugalle, N. R., de Bruyn, L. A. L., & Entwistle, P. (1997). Residual effects of tillage and crop rotation on soil properties, soil invertebrate numbers and nutrient uptake in an irrigated Vertisol sown to cotton. *Applied Soil Ecology*, 7: 11-30.
- Kibar, M., Deniz, K., & Sarıoglu, F. (2012). The morphology, mineralogy, geochemistry and physical implications of foid bearing syenite and syenite-carbonate rocks contact zone soils: Kırşehir-Akpınar-Buzlukdag, Turkey. *Eurasian Journal of Soil Science*, 1(2), 69–74.
- Mzuku, M., Khosla, R., Reich, R., Inman, D., Smith, F., & MacDonald, L. (2005). Spatial variability of measured soil properties across site-specific management zones. *Soil Science Society of America Journal*, 69,(5),1572–1579.
- Ogunkunle, A. O. (1993). Variation of some soil properties along two toposequence on quartzite schist and banded gneiss in South-western Nigeria. *Geo Journal*, 30(4), 399-402.
- Shan, H. M., Liang, H. C., Peng, S. X., Longe, A. A., & Zhou, A. G. (2010). Effects of water-saturation and water-loss processes on composition and structure variations of landslide, Three Gorges reservoir, China. In: *Water- Rock Interaction*. Birkle, P., Torres-Alvarado, I. S. (Eds.). CRC Press, New York, USA.
- Soil Survey Laboratory (2004). *Soil Survey Laboratory Methods Manual*. USDA Natural Resources Conservation Service. Soil Survey Investigations Report No 42. Washington D.C., USA.
- Soil Survey Staff. (2014). *Keys to soil taxonomy*, 12th ed. *USDA-Natural Resources Conservation Service, Washington, DC*.
- Webster, R. (2000). Is soil variation random? *Geoderma*. Volume 97, Issues 3–4, Pages 149-163.
- Webster, R. (2021). Statistical inferential testing. In: Daya Sagar B., Cheng Q., McKinley J., Agterberg F. (eds) *Encyclopedia of Mathematical Geosciences*. *Encyclopedia of Earth Sciences Series*. Springer, Cham.